

LA-UR-18-27446

Approved for public release; distribution is unlimited.

Title: The Geochemistry of Nuclear Fallout

Author(s): Bonamici, Chloe Elizabeth

Kinman, William Scott

Zimmer, Mindy M

Pollington, Anthony Douglas

Rector, Kirk D. Williamson, Todd L. Clifton, Damon Fournelle, John H Hervig, Richard L.

Intended for: Invited talk at the University of Wisconsin Geoscience Department

Issued: 2018-08-06





The Geochemistry of Nuclear Fallout

Chloë Bonamici

LANL

William Kinman

Mindy Zimmer

Anthony Pollington

Kirk Rector

Todd Williamson

Travis Tenner

8 March 2018

<u>UW-Madison</u> John Fournelle


Arizona State Univ Richard Hervig




A Geochemist's Skill Set





1. Thermodynamics & Kinetics






2. Earth Materials Processes






3. Analytical Methods & Instrumentation



Trinitite









Trinitite: What is it Good For?



- Accessible example of nuclear explosion products
- (Cryptic) chemical and isotopic representative of an atomic bomb

Ability to attribute a particular device to a particular source is a nuclear deterrent




Trinity Test

 ²³⁹Pu fuel packed in natural U

Detonated from a steel tower

- Arkosic sand ground material
 - Major minerals: quartz, microcline, anorthoclase, calcite
 - Minor minerals: gypsum, clay, hornblende, salts, zircon






UNCLAS

Trinity Site Geology








Nuclear Fireball Phenomenology






Existing Fallout Formation Model




- Volatilization of device
- Partial melting and entrainment of ground material
- Condensation
 - Refractory bomb material
 - Ground material nuclei
- Homogenization by diffusion and/or melt mixing
- Cooling and fallout





Fallout Heterogeneity

















Trinity Bead Samples






- Element mapping (SEM)
- Major Element Analysis (EPMA)
- Activity mapping (autoradiography)



Major Element Geochemistry






- Total alkali silica diagram
- Compositional diversity beyond known magmatic range



Major Element Geochemistry






- Alkali glass = melted alkali feldspar
- Extreme compositional variability in CaMgFe glass



Major Element Geochemistry






- Overall depletion of volatile elements
- Enrichment in intermediate-refractory elements



Where is the Radioactivity?







 Activity is strongly correlated with regions of CaMgFe glass, consistent with co-condensation of fissionogenic and nonfissionogenic elements



Viscosity & Melt Mixing








 Viscosity contrast likely limited physical mixing of CaMgFe, alkali, and silica melts



Diffusion & Timescales of Cooling








New Fallout Formation Model






- Vaporization of device, tower, sand
- Molten and solid sand entrainment
- Co-condensation of device and ground constituents
- Agglomeration
 - Cooling and fallout



Condensation Chronology & Volatility Scale






 Define a volatility scale based on the relative amounts of volatile and refractory major elements



Fissionogenic Nuclides







 Yields of fissionogenic nuclides are related to the type of fuel used and the energy spectrum of neutrons produced

National Nuclear Security Administration
Slide 20

Trace Elements & Isotopes Methods Los Alamos







- Cameca IMS 6f SIMS for trace elements
- Cameca IMS 1280 SIMS for U-Pu isotopes




- Similar pattern to major elements
- Enrichment of U in beads relative to ground-crust trinitite









Volatility Index


Coherent increasing or decreasing trends

Greatest scatter in most refractory elements



Volatility Index





 Modeling suggests volatile elements experienced kinetic fractionation, whereas refractory elements experienced largely equilibrium fractionation




- Track element fractionation through ratios as a function of volatility index
- Strong element fractionations → sloped trends (e.g., Rb/Sr)
- Weak element fractionations → flat trends (e.g., Zr/Hf)

10³ 10² 10¹ **Frace Element Ratio** 10⁰ 10^{-1} 10⁻² 10^{-3} 10⁻³ 10⁻² 10⁻¹ Volatility Index

Comparison with Freiling (1961)






Stable element-based volatility scales give similar results to fissionogenic isotope-based volatility scales




U-Pu Isotope Geochemistry



 Pu and U fractionated significantly in the Trinity fireball



 U isotopes did not fractionate in the Trinity fireball





Nuclear Forensics Summary



 Test-site sand vaporized and co-condensed along with radioactive bomb material

Glassy fallout preserves compositional variations of the nuclear fireball

- Stable elements are appropriate proxies for studying radionuclide fractionation during nuclear explosions
- Heavy isotopes were not fractionated during the Trinity explosion

NATS AMERICAN PROPERTY Administration


Natural Glasses





Natural Glasses







 Naturally formed glasses show a more limited compositional range than trinitite




Dusty Plasmas





NASA/JPL



Yukawa ball Kading & Melzer 2006



Geochemistry Summary



- Naturally formed glasses show a more limited compositional range than trinitite
- Melting and vaporization behavior of natural glasses differed from those of trinitite, reflecting more heterogeneous energy distributions or greater physical fractionation

 Highly refractory fallout glass may be a process analogue for early solar system condensates like CAIs



Questions?






- This work was supported by:
 - A Los Alamos National Lab
 Seaborg-Strategic Outcomes
 Office Postdoctoral Fellowship to
 C. Bonamici
 - Office of Defense Nuclear
 Nonproliferation R&D Program
 (NA-22), through W. Kinman



U-Pu Isotope Measurements





Cameca IMS 1280

- Static multicollector
- ²³⁴U, ²³⁵U, ²³⁶U, ²³⁸U, 239
- O-primary beam
- 420 cycles, 32 min
- 2200 MRP
- NIST 611, NIST 612
- Hydride correction for ²³⁸U¹H on ²³⁹Pu
 - Ingrowth correction for $^{239}P_{U} \rightarrow ^{235}U$




Oxidation State in the Fireball

Intensity (CCD cts) CH₂/CH₃ bend (1300 & 1430 cm⁻¹) CH_/CH₃ stretch (2850-3030 cm⁻¹)

Raman Shift (cm-1)

Laser Raman spectroscopy

More Diffusion Modeling



Freiling 1961



